

**Supplementary Table 10.** Relative survival rates of gestational trophoblastic neoplasia by age group, SEER stage, and year of diagnosis from 1999 to 2019 in Korea

Variable	N	Relative survival rate (%)					
		3-year	95% CI	5-year	95% CI	10-year	95% CI
Total	1,011	91.8	89.9-93.4	90.5	88.4-92.2	89.8	87.6-91.6
Age group (yr)							
<30	251	95.3	91.7-97.3	94.1	90.2-96.5	93.4	89.4-96.0
30-39	406	92.7	89.6-94.9	91.2	87.8-93.6	90.2	86.7-93.0
40-49	252	89.2	84.5-92.5	88.6	83.8-92.0	87.8	82.7-91.6
≥50	102	86.6	78.1-92.1	83.7	74.5-90.0	83.6	73.9-90.3
Stage at diagnosis (since 2005)							
Localized	249	96.9	93.7-98.6	97.1	93.9-98.8	95.2	90.9-97.7
Regional	47	98.2	86.1-100.0	98.4	86.3-100.2	99.1	86.9-101.0
Distant	224	85.2	79.7-89.3	83.3	77.5-87.7	82.5	76.5-87.2
Unknown	114	94.1	87.7-97.3	92.4	85.5-96.2	93.0	86.0-96.9
Year of diagnosis							
1999-2005	421	90.3	87.0-92.8	88.3	84.8-91.1	87.6	83.9-90.5
2006-2012	362	92.5	89.3-94.9	91.9	88.5-94.4	91.0	87.4-93.7
2013-2019	228	93.9	89.7-96.4	92.6	87.8-95.6		

SEER, Surveillance, Epidemiology, and End Results; CI, confidence interval.