

Supplementary Table 2. Treatment methods applied for vulvovaginitis

Treatment	Total (n=195)	Premenarcheal (n=95)	Postmenarcheal (n=100)	P-value
Observation	110 (56.4)	57 (60.0)	53 (53.0)	0.40
Antiseptics (betadine sitz bath)	59 (30.3)	42 (44.2)	17 (17.0)	<0.01
Local analgesia	7 (3.6)	2 (2.1)	5 (5.0)	0.48
Steroid cream	6 (3.1)	2 (2.1)	4 (4.0)	
Lidocaine cream	1 (0.5)	0 (0.0)	1 (1.0)	
Topical antibiotic ointments	17 (8.7)	13 (13.7)	4 (4.0)	0.03
Antimicrobials	42 (21.5)	15 (15.8)	27 (27.0)	0.08
1st generation cephalosporins	6 (3.1)	5 (5.3)	1 (1.0)	0.19
Cefradroxil	4 (2.1)	3 (3.2)	1 (1.0)	
Cefaclor	2 (1.0)	2 (2.1)	0 (0.0)	
3rd generation cephalosporins	16 (8.1)	6 (6.5)	10 (10.0)	0.50
Cefpodoxim	5 (2.6)	3 (3.2)	2 (2.0)	
Ceftriaxone	3 (1.5)	0 (0.0)	3 (3.0)	
Cefotetan	3 (1.5)	0 (0.0)	3 (3.0)	
Cefdinir	3 (1.5)	1 (1.1)	2 (2.0)	
Cefotaxime	1 (0.5)	1 (1.1)	0 (0.0)	
Cefixime	1 (0.5)	1 (1.1)	0 (0.0)	
Nitroimidazoles (metronidazole)	11 (5.6)	0 (0.0)	11 (11.0)	<0.01
Tetracyclines (doxycycline)	7 (3.6)	0 (0.0)	7 (7.0)	0.03
Penicillin	4 (2.1)	2 (2.1)	2 (2.0)	1.0
Amoxicillin: clavulanate	3 (1.5)	2 (2.1)	1 (1.0)	
Ampicillin	1 (0.5)	0 (0.0)	1 (1.0)	
Sulfonamides	3 (1.5)	1 (1.1)	2 (2.0)	1.0
Macrolides (azithromycin)	2 (1.0)	1 (1.1)	1 (1.0)	1.0
Quinolone (ciprofloxacin)	2 (1.0)	0 (0.0)	2 (2.0)	0.5
Piperacillin-tazobactam	2 (1.0)	0 (0.0)	2 (2.0)	0.5
Carbapenems	1 (0.5)	0 (0.0)	1 (1.0)	1.0
Antifungals	26 (13.4)	6 (6.3)	20 (20.0)	0.01
Clotrimazole	22 (11.3)	6 (6.3)	16 (16.0)	
Fluconazole	4 (2.1)	0 (0.0)	4 (4.0)	

Values are presented as number (%).